

The SAS System

01/11/2017

Site	Monthly RAN goal	RAN goal	SCR visits	BAS visits	RAN visits	SCR:RAN ratio	% of goal	1/2017 SCR visits	1/2017 BAS visits	1/2017 RAN visits	1/2017 SCR:RAN ratio	1/2017 RAN goal(%)	12/2016 SCR visits	12/2016 BAS visits	12/2016 RAN visits	12/2016 SCR:RAN ratio	12/2016 RAN goal(%)
Maine	5.2	125	370	172	107	3.5	85.6	1	.	.	.	.	2	2	2	1.0	38
Tufts	4.7	112	510	262	153	3.3	136.6	1	1	.	.	.	4	6	5	.8	106
MedStar	6.3	150	418	247	159	2.6	106.0	1	1	1	1.0	16	.	.	3	.	48
Duke	6.3	150	375	228	140	2.7	93.3	.	.	.	.	.	3	.	3	1.0	48
SouthCarolina	6.3	150	183	70	32	5.7	21.3	.	.	.	.	.	.	.	.	.	.
Florida	6.3	150	362	122	76	4.8	50.7	.	.	.	.	.	3	.	1	3.0	16
Atlanta	7.5	180	263	227	137	1.9	76.1	.	.	.	.	.	5	4	3	1.7	40
NewYork	4.2	100	190	73	37	5.1	37.0	.	.	.	.	.	.	.	.	.	.
Northwestern	5.2	125	307	220	120	2.6	96.0	.	.	1	.	19	.	.	.	.	.
Tennessee	6.3	150	82	38	29	2.8	19.3	.	.	.	.	.	.	.	.	.	.
Tulane	6.3	150	35	2	2	17.5	1.3	.	.	.	.	.	.	.	.	.	.
Pennington	7.9	190	262	86	47	5.6	24.7	1	1	.	.	.	4	1	.	.	.
HealthPartners	12.5	300	639	474	318	2.0	106.0	.	.	1	.	8	6	5	11	.5	88
Kansas	6.3	150	415	198	121	3.4	80.7	1	3	.	.	.	14	6	3	4.7	48
Baylor	6.3	150	209	82	43	4.9	28.7	.	.	.	.	.	.	.	.	.	.
Nebraska	5.3	126	559	296	175	3.2	138.9	.	.	.	.	.	1	1	4	.3	75
UTSouthwestern	5.0	120	118	78	42	2.8	35.0	.	1	.	.	.	.	.	.	.	.
LA	5.0	120	260	143	84	3.1	70.0	.	.	.	.	.	.	.	.	.	.
Stanford	6.3	150	328	158	91	3.6	60.7	.	.	.	.	.	1	2	2	.5	32
Baltimore	7.7	100	214	125	80	2.7	80.0	.	1	1	.	13	.	4	1	.	13
Cleveland	7.7	100	189	95	55	3.4	55.0	.	.	.	.	.	3	1	1	3.0	13
Kaiser	10.0	130	457	413	265	1.7	203.8	1	1	1	1.0	10	4	1	5	.8	50
Denver	5.4	65	338	136	79	4.3	121.5	.	2	.	.	.	5	.	2	2.5	37
Lenox	5.0	25	51	18	12	4.3	48.0	1	2	.	.	.	7	6	4	1.8	80
	.	.	7134	3963	2404	3.0	100.9	7	13	5	1.4	4	62	39	50	1.2	36

7.1.a Detail on diabetes diagnosis

As of 01/04/17

	Overall	M06	M12	M18	M24	M30	M36	M42	M48
Semi-annual visits	3209	1928	-	1004 ¹	-	277	-	-	-
Diabetes, n (% visits)	82 (2.6)	55 (2.9)	-	19 (1.9)	-	8 (2.9)	-	-	-
Positive A1c and FPG	28	19	-	7	-	2	-	-	-
Positive A1c or FPG, pos. confirmatory	54	36	-	12	-	6	-	-	-
No diabetes, n	3127	1873	-	985	-	269	-	-	-
Negative A1c and FPG	3044	1816	-	967	-	261	-	-	-
Positive A1c or FPG, neg. confirmatory	83	57	-	18	-	8	-	-	-
Weight change (% from baseline)		-0.21%		-0.17%		-0.48%			
Annual visits	2095	-	1437 ¹	-	631	-	27	-	-
Diabetes, n (% visits)	112 (5.3)	-	80 (5.6)	-	29 (4.6)	-	3 (11.1)	-	-
Positive, met 2 criteria	43	-	31	-	10	-	2	-	-
Positive, met 1 criterion, pos. confirmatory	69	-	49	-	19	-	1	-	-
No diabetes, n	1983	-	1357	-	602	-	24	-	-
All 3 criteria negative	1863	-	1272	-	571	-	20	-	-
Met 1 criterion, neg. confirmatory	120	-	85	-	31	-	4	-	-
Weight change (% from baseline)			-0.09%		-0.18%		0.24%		
Outside of study events by adjudication ²	22	4	8	6	2	1	1		
Total events	216	59	88	25	31	9	4		


¹ Number excludes participants who have reached the *diabetes outcome* [by D2d criteria or adjudication].

² If diabetes outcome was adjudicated before stated interval, it is shown under stated interval. For example, adjudicated outcomes between the M06 and M12 visits are shown under M12. Longitudinal analyses (when conducted) will be based on “time-to-event.”


Missed encounters **within last 6 months** (visits and interim phone contact)


3

Study-wide = 5.8%




Inactive and Withdrawn Participants





Discontinuation of Study Pills

excludes participants who withdrew consent



Discontinuation of Study Pills - Last 6 Months

excludes participants who withdrew consent

