






Racial differences in measures of glycemia in the Vitamin D and Type 2 Diabetes (D2d) Study: a secondary analysis of a randomized trial

Erin S LeBlanc ¹, Anastassios G Pittas ², Jason Nelson,³ Raneer Chatterjee,⁴ Neda Rasouli ^{5,6}, Mary K Rhee,⁷ Richard E Pratley ⁸, Cyrus V Desouza ⁹, Lisa M Neff,¹⁰ Anne M Peters,¹¹ Samuel Dagogo-Jack ¹², Daniel S Hsia ¹³, the D2d Research Group

To cite: LeBlanc ES, Pittas AG, Nelson J, *et al*. Racial differences in measures of glycemia in the Vitamin D and Type 2 Diabetes (D2d) Study: a secondary analysis of a randomized trial. *BMJ Open Diab Res Care* 2024;**12**:e003613. doi:10.1136/bmjdr-2023-003613

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/bmjdr-2023-003613>).

Received 14 July 2023
Accepted 12 January 2024



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Erin S LeBlanc;
Erin.S.LeBlanc@kpchr.org

ABSTRACT

Introduction Understanding how race may influence the association between A1c and glycemia can improve diabetes screening. We sought to determine whether, for a given A1c level, glucose levels during an oral glucose tolerance test (OGTT) differed by race.

Research design and methods From data collected at 22 US clinical sites, we conducted a cross-sectional study of concurrently measured A1c and OGTT and observational longitudinal follow-up of the subset with high-risk pre-diabetes. Numerical integration methods were used to calculate area under the glycemic curve (AUC_{glu}) during OGTT and least squares regression model to estimate A1c for a given AUC_{glu} by race, controlling for potential confounders.

Results 1016 black, 2658 white, and 193 Asian persons at risk of diabetes were included in cross-sectional analysis. Of these, 2154 with high-risk pre-diabetes were followed for 2.5 years. For a given A1c level, AUC_{glu} was lower in black versus white participants. After adjustment for potential confounders, A1c levels for a given AUC_{glu} quintile were 0.15–0.20 and 0.02–0.19 percentage points higher in black and Asian compared with white participants, respectively (p<0.05). In longitudinal analyses, black participants were more likely to be diagnosed with diabetes by A1c than white participants (28% vs 10%, respectively; p<0.01). Black and Asian participants were less likely to be diagnosed by fasting glucose than white participants (16% vs 15% vs 37%, respectively; p<0.05). Black participants with A1c levels in the lower-level quintiles had greater increase in A1c over time compared with white participants.

Conclusions Use of additional testing beyond A1c to screen for diabetes may better stratify diabetes risk in the diverse US population.

Hemoglobin A1c (A1c), fasting plasma glucose (FPG), and 2-hour post-load glucose (2hPG) can all be used to diagnose pre-diabetes and diabetes. The relationship of A1c with FPG and the glucose peak and area under the curve during an oral glucose tolerance test (OGTT) was established in

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Previous studies have found that the relationships between glucose and A1c vary between different racial and ethnic groups. However, whether this impacts diabetes diagnosis has not been fully evaluated.

WHAT THIS STUDY ADDS

⇒ We found that A1c levels for a given area under the glycemic curve were higher in black and Asian compared with white participants and that, during follow-up, the test used to diagnose type 2 diabetes varied by race.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ These findings suggest that use of additional testing beyond A1c to screen for diabetes may better stratify diabetes risk in the diverse US population.

the 1970s. Twenty years later, the Diabetes Control and Complications Trial and the UK Prospective Diabetes Study reported that A1c levels were associated with diabetic microvascular complications.^{1 2} These findings led to the adoption of A1c as a method of diagnosing diabetes and pre-diabetes by American Diabetes Association (ADA) and WHO guidelines in 2010 and 2011.^{3 4} Compared with FPG and 2hPG, A1c is unaffected by factors such as exercise and acute stress, does not require fasting or glucose drink consumption, and reflects longer-term glycemic exposure.⁵ A1c also has greater pre-analytical stability, less biological variability, and international standardization.⁵

However, previous studies have found that the relationships between FPG, 2hPG, and A1c vary from person to person and between different racial and ethnic groups. In

particular, A1c has been shown to be 0.1–0.6 percentage points higher in black compared with white individuals for the same degree of glycemia as measured by OGTT, FPG, continuous glucose monitoring, and self-monitored plasma glucose profiles.^{6–11} Differences in the association between A1c and glycemia among people of different races can have an impact on diabetes diagnosis, and this has not been fully evaluated.

The Vitamin D and Type 2 Diabetes (D2d) Study, a diabetes prevention trial conducted from October 2013 to November 2018, included a range of participants by age, gender, race, ethnicity, and spectrum of glycemia (normal glucose regulation, pre-diabetes, diabetes).¹² We collected concurrent data on A1c and glucose measurements during an OGTT, allowing us to examine associations between A1c and both fasting and post-glucose load glycemia. In addition, trial participants underwent repeat glucose testing during follow-up, allowing examination of the potential impact of race on diagnostic testing for diabetes. The D2d cohort, which also collected data on social determinants of health, offers a unique opportunity to examine associations between measures of glycemia by race and ethnicity cross-sectionally and longitudinally.

We hypothesized that for a given area under the glycemic curve (AUC_{glu}) level, A1c would be higher in black individuals compared with individuals who identified as white or Asian, even after controlling for potential confounders. We also hypothesized that the association would remain stable over time and would lead to more black participants being diagnosed with diabetes based on the standard A1c criterion compared with white and Asian participants. We hypothesized that there would be no difference by ethnicity.

RESEARCH DESIGN AND METHODS

Overview of the D2d Study

D2d was a multicenter, randomized, double-blind, placebo-controlled primary prevention clinical trial comparing the effect of 4000 IU of vitamin D₃ versus matching placebo, randomized in a 1:1 ratio, on the incidence of diabetes in people at high risk of diabetes, followed over a median of 2.5 years (ClinicalTrials.gov NCT01942694). The design, including eligibility criteria, when and where data were collected, a description of the intervention, and how randomization was implemented, and main results of D2d have been published.^{12 13}

Baseline characteristics, including social determinants of health and vitamin D level

Self-administered questionnaires were used to collect data on age, gender, smoking status (never, former, current), annual household income (<\$75 000 vs ≥\$75 000), and education (no high school, no post-high school, some post-high school, bachelor's degree, graduate degree). Other self-administered questionnaires included the International Physical Activity Questionnaire¹⁴ to determine total physical activity per week

(metabolic equivalents-hours/week) and the Multi-cultural Food Frequency Questionnaire to determine alcohol use (g/day) and dietary glycemic index. Body mass index (BMI) (kg/m^2) was calculated using weight and height measured using standardized procedures. The 25-hydroxyvitamin D level (25(OH)D) was measured by liquid chromatography–tandem mass spectrometry, as previously described.¹²

Race and ethnicity

Reporting race and ethnicity in this study was mandated by the US National Institutes of Health (NIH). Race and ethnicity were self-reported by study participants. Per NIH guidance, race was categorized as American Indian or Alaska Native, Asian, black or African American, Native Hawaiian or Other Pacific Islander, white, or more than one race. Due to the low number of participants in certain racial categories, our analytical sample for the analysis by race was restricted to participants who self-identified their race as black, white, or Asian (online supplemental figure 1). Ethnicity was categorized as Hispanic or not Hispanic.

Glycemic testing

D2d used a two-step screening process (online supplemental figure 1). At screening step 1, FPG and A1c were measured. If results suggested participants were likely to meet D2d glycemic enrollment criteria, they were invited for a full screening visit. At the full screening visit, which also served as the baseline visit for participants who were randomized, participants underwent a 75-gram OGTT to measure FPG, glucose at 30 min (Glu-30) and 2hPG, along with A1c. Participants were instructed not to change physical activity and diet for the 3 days before the OGTT. All tests were analyzed by the D2d central laboratory to determine final eligibility for the trial. A1c was measured by an ion-exchange high-performance liquid chromatography method certified by the National Glycohemoglobin Standardization Program.¹⁵ Plasma glucose was measured with the use of a hexokinase method (Roche Cobas Integra at the University of Vermont Laboratory for Clinical Biochemistry Research).¹²

Participants

Cross-sectional

The cross-sectional analyses include participants who underwent glycemic testing at the full screening visit (online supplemental figure 1; $n=3876$), regardless of their randomization status in the trial.

Longitudinal

Eligibility criteria for D2d included meeting at least two of the glycemic criteria for pre-diabetes per the 2010 ADA guidelines: FPG 100–125 mg/dL (5.6–6.9 mmol/L), 2hPG 140–199 mg/dL (7.8–11.0 mmol/L), or A1c 5.7–6.4% (39–46 mmol/mol).^{16 17} Participants were also required to be age ≥30 years (≥25 years for American Indians, Alaska Natives, Native Hawaiians, or other Pacific Islanders) and have a BMI 24–42 kg/m^2 (22.5–42 kg/m^2)

for Asian). Exclusion criteria included use of medications approved for treatment of diabetes, history of bariatric surgery, chronic kidney disease (estimated glomerular filtration rate $<50\text{ mL/min}/1.73\text{ m}^2$)¹³ and presence of hemoglobin variants identified during screening that are known to interfere with the A1c assay, such as hemoglobin E, but not hemoglobin S or hemoglobin C.¹⁸

Participants who were randomized in D2d and self-reported to be of black, white or Asian race are included in the present longitudinal analyses if they had at least one follow-up glycemic measure (online supplemental figure 1; $n=2154$). Glycemic status was assessed annually with FPG, A1c and 2hPG, and semiannually with FPG and A1c. During the trial, if at least two of the glycemic measures met the ADA thresholds for diabetes,¹⁶ the participant was considered to have met the diabetes outcome. When only one glycemic measure met the threshold, confirmatory testing was performed. A diagnosis of diabetes made outside of D2d that was not confirmed by the D2d central laboratory testing was not considered in the present analysis.

Derived glycemic values

To measure the glucose excursion during an OGTT, we used optimum numerical integration methods to estimate AUC_{glu} from FPG and plasma glucose at 30 min and at 2 hours.¹⁹ We then categorized participants into AUC_{glu} quintiles. The integration of 3 glucose points provides a better picture of the overall glycemia versus examining individual time points.

Statistical analysis

Baseline difference in potential confounders

Descriptive statistics included percentage, means \pm SD, or medians (IQR: Q1–Q3) for non-normally distributed data. Comparisons between racial and ethnic groups at baseline used Fisher's exact test, the χ^2 test, the Wilcoxon rank-sum test, or the pooled-variance t-test. Given the large Ns in our population, we used standard mean difference (SMD) to examine the strength of the association with racial and ethnic categories (>0.2 or <-0.2 indicates a moderate association; $\text{SMD} >0.5$ or <-0.5 indicates a large association).

Cross-sectional analysis

We examined the relationship between A1c and AUC_{glu} and component glucose measures (FPG, Glu-30 and 2hPG) stratified by self-identified race (black, white, or Asian). The linear intercept and slope with 95% CIs between glucose measures and A1c were plotted for each race. Linearity was evaluated by adding higher-order polynomial terms to the model and considered statistically significant at $p < 0.05$. We repeated the analysis using ethnicity instead of race. In addition, we repeated the race analyses excluding those with hemoglobin variants. The primary analysis was a least squares regression model with A1c as the dependent variable and AUC_{glu} , race, and the interaction between them as independent variables. All

reported p values were two-sided. To account for potential confounders in the association between A1c and AUC_{glu} , we then estimated racial and ethnic mean differences and 95% CIs in A1c for a given AUC_{glu} , controlling for the following potential confounders: age, gender, BMI, the interaction between age and gender and BMI, alcohol use, education, income, presence of hemoglobin variants, smoking status, physical activity, dietary glycemic index, and baseline 25(OH)D level; we refer to this as adjusted A1c.

Longitudinal analysis

We tested whether the distribution of diabetes diagnoses differed by race and ethnicity using χ^2 tests of association. We also examined the change in A1c over time in black and white participants. We estimated the average per cent change from baseline (and 95% CIs) of A1c in each AUC_{glu} quintile for each race and the difference between race groups using linear mixed-effects regression models to account for repeated measurements over time. The unadjusted and adjusted models used A1c as the dependent variable and AUC_{glu} , baseline A1c, race, and the interaction between race and AUC_{glu} as independent variables. All reported p values were two-sided. The adjusted model controlled for the same potential confounders as the primary analysis.

RESULTS

Baseline characteristics

The cross-sectional analysis included 1016 black, 2658 white, and 193 Asian participants with complete glycemic measurements (FPG, Glu-30, 2hPG), of whom 323 were Hispanic and 3544 were non-Hispanic. Their baseline characteristics are shown in online supplemental tables 1 and 2. White participants were older, had higher 25(OH)D levels, and consumed more alcohol than black and Asian participants. Black participants had lower hemoglobin levels, were more likely to have hemoglobin S trait, and were less likely to earn \$75 000 or more per year or have a graduate or professional degree than white participants. Asian participants had lower BMIs, higher incomes, higher education levels, and lower physical activity levels than white participants. Compared with white participants, a larger proportion of black participants had an A1c in the pre-diabetes range and a smaller proportion had an FPG in the pre-diabetes range. Black participants were more likely than white participants to have all three tests in the normal range. Hispanic participants were younger than non-Hispanic participants (online supplemental table 2), relatively more women self-identified as Hispanic than non-Hispanic, and Hispanic participants had lower incomes and education levels than non-Hispanic participants.

The longitudinal analysis included 2359 participants (616 black, 1613 white, and 130 Asian) who were randomized and followed over time: of whom, 190 were Hispanic and 2169 were non-Hispanic. Differences in baseline




Figure 1 Cross-sectional association* between A1c and AUC_{glu} and fasting, 30-minute, and 2-hour plasma glucose by race. (A) Association between A1c and AUC_{glu} in black versus white participants; (B) association between A1c and fasting glucose in black versus white participants; (C) association between A1c and 30-minute glucose in black versus white; (D) association between A1c and 2-hour glucose in black versus white participants; (E) association between A1c and AUC_{glu} in Asian versus white; (F) association between A1c and fasting glucose in Asian versus white participants; (G) association between A1c and 30-minute glucose in Asian versus white participants; (H) association between A1c and 2-hour glucose in Asian versus white participants. *Shaded areas represent 95% CIs of the regression curve. AUC_{glu} , area under the glycemic curve.

characteristics of those randomized in D2d by race were similar to those seen in the screening population (online supplemental tables 3 and 4).

Baseline AUC_{glu} categories

The quintiles of AUC_{glu} for the full screening cohort are shown in online supplemental figure 2, along with the mean FPG, Glu-30 and 2hPG values for each AUC_{glu} quintile. The mean FPG, Glu-30, and 2hPG level for the first quintile were 98.7, 140.2, and 95.2 mg/dL compared with 115.5, 208.6, and 198.3 mg/dL in the highest quintile, respectively.

Cross-sectional association between A1c and AUC_{glu} , FPG, 2hPG by race

For a given A1c level, the AUC_{glu} was lower in black versus white participants (figure 1A). A similar pattern emerged when we examined FPG, Glu-30, or 2hPG for a given A1c among black and white participants (figure 1B–D). Differences were less pronounced for 2hPG at lower A1c levels. When we excluded the 37 with hemoglobin variants (ie, hemoglobin S and hemoglobin C), the associations did not change (data not shown). When we compared Asian with white participants, overall AUC_{glu} for a given A1c was not different. However, Asian participants did have lower FPG for a given A1c compared with white participants, although there were no differences in AUC_{glu} levels by Glu-30 and 2hPG (figure 1E–H). There were no significant differences in AUC_{glu} for a given A1c

between Hispanic versus non-Hispanic participants (data not shown).

When we estimated racial and ethnic mean differences and 95% CIs in A1c for a given AUC_{glu} controlling for potential confounders, the adjusted A1c levels for a given AUC_{glu} quintile were 0.15–0.20 and 0.02–0.19 percentage points higher in black ($p < 0.01$ all quintiles) and Asian ($p < 0.05$ for quintiles 1, 3–5; NS for quintile 2), compared with white, participants (table 1). In the highest AUC_{glu} quintile, Hispanic participants had a higher adjusted A1c compared with non-Hispanic participants (6.05 (5.98, 6.12) vs 5.96 (5.95, 5.98); $p < 0.03$), but there were no A1c differences between ethnic groups in the other AUC_{glu} quintiles (online supplemental table 5).

Racial differences in tests confirming diagnosis of diabetes during follow-up

When the diagnosis of diabetes was made during study trial follow-up, the diagnostic tests confirming diagnosis differed by race. Of those who developed diabetes during the study, 27.5% of black participants vs 9.8% of white participants were diagnosed by two A1c values ($p < 0.01$; table 2) and 19.2% vs 12.0% were diagnosed by simultaneously elevated A1c and FPG ($p = 0.04$), respectively. A much larger proportion of white participants (37.3%) were diagnosed with diabetes by two FPG values compared with 15.8% and 15.2% of black and Asian participants ($p < 0.01$ and $p = 0.01$, respectively). There were no other differences between racial groups. These

Table 1 A1c level by AUC-glucose quintiles in screening cohort by race

AUC-glucose quintile*	Black		White		Asian	
	Mean (range)	Observed	Adjusted†	Observed	Adjusted†	Observed
Low—1	5.91 (5.89, 5.94)‡	5.94 (5.91, 5.97)‡	5.73 (5.71, 5.75)	5.74 (5.71, 5.76)	5.85 (5.74, 5.95)§	5.87 (5.77, 5.98)‡
2	5.96 (5.93, 5.99)‡	5.98 (5.95, 6.02)‡	5.78 (5.76, 5.81)	5.78 (5.76, 5.80)	5.78 (5.69, 5.86)	5.80 (5.71, 5.89)
3	5.96 (5.93, 6.00)‡	5.98 (5.94, 6.01)‡	5.82 (5.80, 5.84)	5.81 (5.79, 5.83)	5.94 (5.87, 6.01)‡	6.00 (5.92, 6.07)‡
4	6.02 (5.98, 6.06)‡	6.02 (5.98, 6.06)‡	5.86 (5.84, 5.88)	5.84 (5.82, 5.86)	5.91 (5.84, 5.98)	5.93 (5.86, 6.00)§
High—5	6.07 (6.03, 6.12)‡	6.09 (6.04, 6.14)‡	5.95 (5.93, 5.97)	5.94 (5.92, 5.96)	5.85 (5.74, 5.95)	6.03 (5.96, 6.10)§

*Quintiles were derived for the screening population (all races combined).

†Estimated linear regression for adjusted A1c; $R^2=0.046$ for observed model (no adjustment); $R^2=0.119$ for model adjusted for race+race×AUC quintile group; $R^2=0.137$ for model adjusted for race+AUC quintile groups+race×AUC quintile groups+age×gender×BMI; $R^2=0.148$ for fully adjusted model=(race)+AUC quintile groups+(race)×AUC quintile groups+adjusted variables (age, gender, BMI, age×gender×BMI, alcohol use, education, income, presence of hemoglobinopathy, smoking status, physical activity, dietary glycemic index, and 25(OH)D ($R^2=0.148$)).

‡ $P<0.01$ for white versus black participants or white versus Asian participants.

§ $P<0.05$ for white versus Asian participants.

AUC, area under the curve; BMI, body mass index; 25(OH)D, 25-hydroxyvitamin D.

differences are likely attributable to black participants having higher A1c levels at diagnosis compared with white participants (mean (SD): 6.5 (0.6) vs 6.3 (0.5); $p<0.001$). FPG and 2hPG did not differ between racial groups at diabetes diagnosis. Although mean A1c (SD) was higher at diabetes diagnosis in Hispanic versus non-Hispanic participants (6.5 (0.6) vs 6.3 (0.5); $p=0.02$), FPG and 2hPG were not different between ethnic groups at diabetes diagnosis (data not shown).

Longitudinal change in A1c over time by baseline AUC_{glu} quintile by race

Average change in A1c compared with baseline did not differ by treatment group and so the vitamin D and

placebo groups were combined in the longitudinal analysis. The absolute increase in A1c over time according to baseline AUC_{glu} quintiles differed by race (online supplemental table 6). Compared with white participants, black participants in lower baseline AUC_{glu} quintiles had greater increases in A1c over time (first quintile: 0.76 (0.29, 1.23) vs 0.03 (−0.37, 0.43); unadjusted $p=0.02$ and second quintile: 1.81 (1.23, 2.40) vs 0.85 (0.48, 1.21); unadjusted $p=0.006$, respectively). When adjusted for potential confounders, the p value for A1c change in the first quintile was no longer significant (adjusted $p=0.057$) but remained significant for those in the second quintile (adjusted $p=0.027$). A1c increased similarly in white and

Table 2 Glycemic tests at diabetes diagnosis by race in enrolled population (longitudinal analysis)

Observed level, mean (SD)	Overall n=563	Race			P value	White vs	White vs
		Black n=120	White n=410	Asian n=33		black	Asian
Fasting plasma glucose (FPG), mg/dL	127.6 (22.3)	128.7 (31.5)	127.7 (19.5)	121.8 (12.9)	0.28	0.91	0.30
2-hour post-load glucose (2hPG), mg/dL	205.7 (47.9)	215.7 (50.1)	202.3 (45.2)	213.5 (62.8)	0.10	0.12	0.50
Hemoglobin A1c, %	6.3 (0.5)	6.5 (0.6)	6.3 (0.5)	6.4 (0.3)	<0.01	<0.001	0.43
Diabetes confirmatory tests, no (%)					<0.001		
2 A1c tests	77 (13.7)	33 (27.5)	40 (9.8)	4 (12.1)		<0.01	0.66
2 FPG tests	177 (31.4)	19 (15.8)	153 (37.3)	5 (15.2)		<0.01	0.01
2 2hPG tests	109 (19.4)	16 (13.3)	83 (20.2)	10 (30.3)		0.09	0.17
A1c+FPG	77 (13.7)	23 (19.2)	49 (12.0)	5 (15.2)		0.04	0.59
FPG+2hPG	62 (11.0)	12 (10.0)	46 (11.2)	4 (12.1)		0.71	0.87
2hPG+A1c	34 (6.0)	10 (8.3)	21 (5.1)	3 (9.1)		0.19	0.33
A1c+FPG+2hPG	27 (4.8)	7 (5.8)	18 (4.4)	2 (6.1)		0.51	0.66

black persons in the three highest baseline AUC_{glu} quintiles before and after adjustment.

CONCLUSIONS

This study confirms previous findings that black individuals have higher A1c levels than white participants and adds information about the association between glucose and A1c in the less well-studied Asian population. It also extends previous work by relating baseline A1c to incident diabetes and change in A1c over a 2.5-year period according to race.

The present analyses focused on understanding how race may impact the association between A1c and glycemia as measured by AUC_{glu} during an OGTT in a population with a range of glycemia (from normal to diabetes). Most of the prior work has examined the association between A1c and glycemia by race using adjustment of A1c by glucose levels. These studies have generally found that black participants have higher A1c levels compared with white participants.^{6–8 10 20} A different analytic approach involving estimation of A1c for a given glucose level was used by this study, as well as two prior studies.^{8 21} In one of these previous studies, which was conducted in those with type 1 diabetes, black individuals had a higher A1c for a given mean glucose as measured by continuous glucose monitoring (CGM).⁸ In the other study, which consisted predominantly of those with type 1 and type 2 diabetes, association between A1c and mean glucose during fingerstick monitoring was found.²¹ Our study adds to this prior work by estimating racial mean differences and 95% CIs in A1c for a given AUC_{glu} controlling for potential confounders in a population consisting predominantly of those with pre-diabetes. While we found that race had a small but significant association with A1c for a given level of mean glycemia, other factors beyond race may also contribute to explained variance. Because these study data were not specifically collected for this post-hoc analysis, the results are preliminary and can only suggest possible differences.

The association between A1c and glucose has been less well studied in other races and ethnic groups. We noted that Asian participants with a given A1c level had lower FPG levels, but not AUC_{glu}, compared with white participants and after adjustments for potential covariates, particularly BMI, A1c was generally higher for a given AUC_{glu} quintile. Previous work examining A1c levels in Asian compared with white individuals has been mostly in those with diabetes and has had mixed findings.^{7 10 21} Our study adds important data on a possible clinically important difference in the A1c–glucose association among Asian and white individuals with pre-diabetes. Previous work from the National Health and Nutrition Examination Survey found that Mexican-Americans were more likely to have an elevated A1c than non-Hispanic white people.²² In studies of those with high risk of diabetes and established diabetes, A1c levels were higher in Hispanic versus non-Hispanic participants

after adjustment for glucose levels,¹⁰ which matches our finding that adjusted A1c differed by ethnicity in the highest AUC_{glu} quintile.

It has been proposed that racial differences in the glucose–A1c relationship could be due to differences in prevalence of hemoglobin variants,²³ but when we excluded those with abnormal hemoglobin variants, the associations remained unchanged. Differences in how much glucose attaches to hemoglobin for a given blood glucose concentration have also been invoked as an explanation for the differences in associations between A1c and glucose measures by race. Other explanations for the racial and ethnic differences in the A1c–glucose relationship include differences in red blood cell, age and turnover, lower average hemoglobin levels, and genetic differences.^{24–26} There may also be other factors that were not assessed in the present study (including social determinants and gluoregulatory function) that might impact long-term glycemic excursions that may not be captured by measurement of FPG or 2hPG.^{26 27}

The most novel piece of our work is the longitudinal follow-up, which reveals that racial differences in the glucose–A1c relationship likely have a clinical impact. Among black participants, A1c was more likely than FPG or 2hPG to diagnose both high-risk pre-diabetes at baseline and incident diabetes during study follow-up. On the other hand, white participants were more likely to be diagnosed with diabetes using FPG compared with black and Asian participants. Therefore, if A1c alone is used to screen for pre-diabetes and diabetes, A1c differences in black and Asian compared with white individuals would lead to differential diabetes diagnoses. Descriptively, the 2-hour time point during the OGTT appeared a better approximation of A1c in black persons compared with the other concurrent time points. Use of additional testing beyond A1c to screen for diabetes may better stratify diabetes risk in the diverse US population, especially using the 2-hour OGTT which showed less divergence between black and white participants. While the choice of screening tests may influence when individuals in different ethnic groups are diagnosed with diabetes, it is unclear whether these differences in individual's glycemic testing at the time of diabetes diagnosis have long-term impact on the development of microvascular and macrovascular complications.²⁸

We also examined longitudinal change in A1c by baseline AUC_{glu} quintile according to race. Our results suggest that black participants who have the lowest AUC_{glu} during an OGTT (and theoretically the least risk of progression to diabetes) have a more rapid rise of A1c than white participants who are in the same low AUC_{glu} category. This finding needs confirmation in other cohorts to better understand how frequently A1c needs to be monitored among the diverse US population who are in the early stages of hyperglycemia.

Our study has several strengths, including a large sample size, extensive phenotyping of people at risk of diabetes, and longitudinal follow-up from pre-diabetes

to diabetes. Our study also had some limitations. The screening cohort for the cross-sectional analysis had already been through a pre-screening process which likely resulted in a screened population at relatively higher risk than the general population. We were also limited by only having a single OGTT per person, while A1c reflects approximately 2 months of glycemia. Future research should examine racial differences in fasting and postprandial glucose levels over longer periods of time, with multiple test results and/or CGM, reflecting the same time window as A1c.

In conclusion, our work confirms previous studies reporting that A1c is higher for a given mean glucose level in those who self-identify as black compared with white even after adjustment for potential covariates. This finding indicates that choice of screening tests for pre-diabetes and diabetes may lead to different diagnostic outcomes in different racial groups, particularly between white and black individuals. Such differences in mean glycemia at the time of diabetes diagnosis may have important clinical implications with regard to how early diabetes is detected which can have an impact on management and development of long-term complications; these questions require additional study.

Author affiliations

¹Kaiser Permanente Center for Health Research, Portland, Oregon, USA

²Division of Endocrinology, Diabetes and Metabolism, Tufts Medical Center, Boston, Massachusetts, USA

³Tufts Medical Center, Boston, Massachusetts, USA

⁴Internal Medicine, Duke University, Durham, North Carolina, USA

⁵Endocrinology, Metabolism and Diabetes, University of Colorado Denver School of Medicine, Aurora, Colorado, USA

⁶Endocrinology, VA Eastern Colorado Health Care System, Denver, Colorado, USA

⁷Medicine/Endocrinology, Metabolism, and Lipids, Emory University School of Medicine, Atlanta, Georgia, USA

⁸Translational Research Institute, AdventHealth Research Institute, Orlando, Florida, USA

⁹University of Nebraska Medical Center, Omaha, Nebraska, USA

¹⁰Northwestern Medicine, Chicago, Illinois, USA

¹¹Endocrinology, USC, Manhattan Beach, California, USA

¹²Division of Endocrinology, Diabetes & Metabolism General Clinical Research Center, University of Tennessee Health Science Center, Memphis, Tennessee, USA

¹³Pennington Biomedical Research Center, Baton Rouge, Louisiana, USA

Acknowledgements The authors thank the D2d investigators, staff, and trial participants for their outstanding dedication and commitment to the study. AGP is supported in part by generous donations to the Tupper Research Fund at Tufts Medical Center.

Collaborators D2d Research Group collaborators listed in the online supplemental file. Steering Committee: Anastassios G. Pittas, MD MS, Tufts Medical Center, Boston, MA (Chair); Irwin Brodsky, MD, Maine Medical Center Research Institute, Scarborough, ME; Lisa Ceglia, MD MS, Tufts Medical Center, Boston, MA; Chhavi Chadha, MD, HealthPartners Research Foundation, Minneapolis, MN; Rane Chatterjee, MD MPH, Duke University Medical Center, Durham, NC; Bess Dawson-Hughes, MD, Jean Mayer USDA Human Nutrition Research Center on Aging at Tufts University, Boston, MA; Cyrus Desouza, MBBS, Omaha VA Medical Center, University of Nebraska Medical Center, Omaha, NE; Rowena Dolor, MD MHS, Duke University Medical Center, Durham, NC; John Foreyt, PhD, Baylor College of Medicine, Houston, TX; Adline Ghazi, MD, MedStar Good Samaritan Hospital, Baltimore, MD; Daniel S. Hsia, MD, Pennington Biomedical Research Center, Baton Rouge, LA; Karen C. Johnson, MD MPH, University of Tennessee Health Science Center, Memphis, TN; Sangeeta R. Kashyap, MD, Cleveland Clinic, Cleveland, OH; Sun Kim, MD, Stanford University Medical Center, Stanford, CA; Erin S. LeBlanc, MD

MPH, Kaiser Permanente Center for Health Research NW, Portland, OR; Michael R. Lewis, MD MBA, University of Vermont—Central Laboratory, Burlington, VT; Emilia Liao, MD, Northwell Health Lenox Hill Hospital, New York, NY; Saul Malozowski, MD PhD, National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda, MD (NIDDK Program Officer); Lisa M. Neff, MD, Northwestern University, Chicago, IL; Patrick O'Neil, PhD, Medical University of South Carolina, Charleston, SC; Jean Park, MD, MedStar Health Research Institute, Hyattsville, MD; Anne Peters, MD, Keck School of Medicine of the University of Southern California, Los Angeles, CA; Lawrence S. Phillips, MD, Atlanta VA Medical Center, Decatur, GA and Emory University School of Medicine, Atlanta, GA; Richard Pratley, MD, AdventHealth Translational Research Institute for Metabolism and Diabetes, Orlando, FL; Philip Raskin, MD, University of Texas Southwestern Medical Center, Dallas, TX; Neda Rasouli, MD, University of Colorado, School of Medicine and VA Eastern Colorado Health Care System, Aurora, CO; David Robbins, MD, University of Kansas Medical Center, Kansas City, KS; Clifford Rosen, MD, Maine Medical Center Research Institute, Scarborough, ME; Past Steering Committee members: Vanita R. Aroda, MD, Brigham and Women's Hospital, Boston, MA; Patricia Sheehan, RN MPH MS, Spaulding Rehabilitation Network, Boston, MA; Myrlene A. Staten, MD, National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda, MD (NIDDK Program Scientist); Advisor: William C. Knowler, MD DrPH, National Institute of Diabetes and Digestive and Kidney Diseases, Phoenix, AZ

Contributors ESL, AGP, JN, RC, NR, MKR, REP, CD, LMN, AP, and DSH were involved in the conception, design, and conduct of the study and the analysis and interpretation of the results. ESL wrote the first draft of the manuscript, and all authors edited, reviewed, and approved the final version of the manuscript. ESL, AGP, and JN are the guarantors of this work and, as such, had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Funding The conduct of the trial was supported primarily by the NIDDK and the Office of Dietary Supplements of the National Institutes of Health through the multicenter clinical study cooperative agreement (U01DK098245; principal investigator, AGP) to Tufts Medical Center, where the D2d Coordinating Center is based. The U01 grant mechanism establishes the NIDDK project scientist (Dr Staten) as a member of the D2d Research Group. The trial also received secondary funding from the American Diabetes Association to Tufts Medical Center (1-14-D2d-01; principal investigator, AGP).

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval This study involves human participants. The Tufts Health Sciences IRB served as IRB of the coordinating center, and each of the 22 participating sites' IRBs also approved the study: Tufts Health Sciences IRB Coordinating Center (10594); Maine Medical Center/MaineHealth IRB (958790-9); Tufts Health Sciences IRB, Clinical Site (10866); MedStar Health Research Institute IRB (2013-130); Duke Health IRB (47517); Medical University of South Carolina IRB for Human Research (24354); Florida Hospital IRB (481944); Emory University IRB (68082); Mt Sinai Program for the Protection of Human Subjects IRB (16-00216); Northwestern University IRB (78718); University of Tennessee Health Science Center IRB (13-02590); Pennington Biomedical Research Center IRB (2013-042-PBRC D2d); HealthPartners Institute IRB (A12-056); University of Kansas Medical Center IRB (13811); Baylor College of Medicine IRB (H-33130); University of Nebraska Medical Center IRB (411-13-FB); UT Southwestern IRB (8843); University of Southern California Health Sciences Campus IRB (13-00450); Stanford University IRB (27848); Omaha VA Medical Center IRB (874); Kaiser Permanente IRB (4927); Cleveland Clinic IRB (14-718); Colorado Multiple IRB (14-2402). All participants provided informed consent.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. Datasets generated and/or analyzed during the current study and the associated data dictionary are not publicly available but are available from the D2d Coordinating Center at Tufts Medical Center on reasonable request by bona fide researchers after acceptance for publication. Protocol synopsis, contact details, publications, and the process for collaboration and data requests can be found on the website (d2dstudy.org).

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability

of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Erin S LeBlanc <http://orcid.org/0000-0002-4067-3432>
 Anastassios G Pittas <http://orcid.org/0000-0002--0124-7174>
 Neda Rasouli <http://orcid.org/0000-0003-1269-3580>
 Richard E Pratley <http://orcid.org/0000-0002-2912-1389>
 Cyrus V Desouza <http://orcid.org/0000-0001-6660-0568>
 Samuel Dagogo-Jack <http://orcid.org/0000-0001-5318-9677>
 Daniel S Hsia <http://orcid.org/0000-0003-2832-3429>

REFERENCES

- Diabetes Control and Complications Trial Research Group, Nathan DM, Genuth S, *et al.* The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med* 1993;329:977–86.
- Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet* 1998;352:837–53.
- American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2010;33 Suppl 1:S62–9.
- World Health Organization. Use of glycated haemoglobin (HbA1c) in diagnosis of diabetes mellitus: abbreviated report of a WHO consultation. Secondary use of glycated haemoglobin (HbA1c) in diagnosis of diabetes mellitus: abbreviated report of a WHO consultation. 2011. Available: <https://iris.who.int/handle/10665/70523>
- Sacks DB. A1C versus glucose testing: a comparison. *Diabetes Care* 2011;34:518–23.
- Ziemer DC, Kolm P, Weintraub WS, *et al.* Glucose-independent, black-white differences in hemoglobin A1c levels: a cross-sectional analysis of 2 studies. *Ann Intern Med* 2010;152:770–7.
- Herman WH, Ma Y, Uwaifo G, *et al.* Differences in A1C by race and ethnicity among patients with impaired glucose tolerance in the diabetes prevention program. *Diabetes Care* 2007;30:2453–7.
- Bergenstal RM, Gal RL, Connor CG, *et al.* Racial differences in the relationship of glucose concentrations and hemoglobin A1c levels. *Ann Intern Med* 2017;167:95–102.
- Lipska KJ, De Rekeneire N, Van Ness PH, *et al.* Identifying dysglycemic states in older adults: implications of the emerging use of hemoglobin A1c. *J Clin Endocrinol Metab* 2010;95:5289–95.
- Rasouli N, Younes N, Utzschneider KM, *et al.* Association of baseline characteristics with insulin sensitivity and β -Cell function in the glycemia reduction approaches in diabetes: a comparative effectiveness (GRADE) study cohort. *Diabetes Care* 2021;44:340–9.
- Herman WH, Dungan KM, Wolffenbuttel BHR, *et al.* Racial and ethnic differences in mean plasma glucose, hemoglobin A1c, and 1,5-anhydroglucitol in over 2000 patients with type 2 diabetes. *J Clin Endocrinol Metab* 2009;94:1689–94.
- Pittas AG, Dawson-Hughes B, Sheehan P, *et al.* Vitamin D supplementation and prevention of type 2 diabetes. *N Engl J Med* 2019;381:520–30.
- Pittas AG, Dawson-Hughes B, Sheehan PR, *et al.* Rationale and design of the Vitamin D and Type 2 Diabetes (D2d) study: a diabetes prevention trial. *Diabetes Care* 2014;37:3227–34.
- Craig CL, Marshall AL, Sjöström M, *et al.* International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 2003;35:1381–95.
- Little RR, Rohlfing C, Sacks DB. The National glycohemoglobin standardization program: over 20 years of improving hemoglobin A_{1c} measurement. *Clin Chem* 2019;65:839–48.
- American Diabetes Association. Standards of medical care in diabetes--2010. *Diabetes Care* 2010;33 Suppl 1:S11–61.
- American Diabetes Association Professional Practice Committee. 2. Classification and diagnosis of diabetes: standards of medical care in diabetes-2022. *Diabetes Care* 2022;45:S17–38.
- Lewis MR, Macauley RC, Sheehan PR, *et al.* Management of hemoglobin variants detected incidentally in HbA1c testing: a common problem currently lacking a standard approach. *Diabetes Care* 2017;40:e8–9.
- Purves RD. Optimum numerical integration methods for estimation of area-under-the-curve (AUC) and area-under-the-moment-curve (AUMC). *J Pharmacokinetics Biopharm* 1992;20:211–26.
- Chapp-Jumbo E, Edeoga C, Wan J, *et al.* Ethnic disparity in hemoglobin A1c levels among normoglycemic offspring of parents with type 2 diabetes mellitus. *Endocr Pract* 2012;18:356–62.
- Nathan DM, Kuenen J, Borg R, *et al.* Translating the A1C assay into estimated average glucose values. *Diabetes Care* 2008;31:1473–8.
- Menke A, Rust KF, Savage PJ, *et al.* Hemoglobin A1c, fasting plasma glucose, and 2-hour plasma glucose distributions in U.S. population subgroups: NHANES 2005-2010. *Ann Epidemiol* 2014;24:83–9.
- Klonoff DC. Hemoglobinopathies and hemoglobin A1c in diabetes mellitus. *J Diabetes Sci Technol* 2020;14:3–7.
- Hivert M-F, Christophi CA, Jablonski KA, *et al.* Genetic ancestry markers and difference in A1c between African American and White in the diabetes prevention program. *J Clin Endocrinol Metab* 2019;104:328–36.
- Wheeler E, Leong A, Liu C-T, *et al.* Impact of common genetic determinants of hemoglobin A1c on type 2 diabetes risk and diagnosis in ancestrally diverse populations: a transethnic genome-wide meta-analysis. *PLoS Med* 2017;14:e1002383.
- Sarnowski C, Leong A, Raffield LM, *et al.* Impact of rare and common genetic variants on diabetes diagnosis by hemoglobin A1c in multi-ancestry cohorts: the trans-omics for precision medicine program. *Am J Hum Genet* 2019;105:706–18.
- Ebenibo S, Edeoga C, Wan J, *et al.* Glucoregulatory function among African Americans and European Americans with normal or pre-diabetic hemoglobin A1c levels. *Metabolism* 2014;63:767–72.
- Gore MO, McQuire DK. A test in context: hemoglobin A1c and cardiovascular disease. *J Am Coll Cardiol* 2016;68:2479–86.



Vitamin **D** and
type **2** diabetes

Racial Differences in Glycemia and Hemoglobin A1c in the Vitamin D and Type 2

Diabetes (D2d) study (First Author: Erin S. LeBlanc, MD)

ONLINE-ONLY SUPPLEMENTAL MATERIAL

D2d Research Group collaborators.....	2
Supplemental Figure S1: Consort diagram	4
Supplemental Table S1: Baseline characteristics of screening cohort by ethnicity (cross-sectional analyses)	7
Supplemental Table S2: Baseline characteristics of enrolled cohort by race (longitudinal analyses)	12
Supplemental Table S3: Baseline characteristics of enrolled cohort by ethnicity	14
Supplemental Table S4: Adjusted A1c level by AUC-glucose quintiles in screened population by ethnicity.....	16
Supplemental Table S5: Change in observed A1c over time by baseline AUC-glucose quintile and race in enrolled cohort (longitudinal)	17



D2d Research Group collaborators

Steering Committee

Anastassios G. Pittas, MD MS, Tufts Medical Center, Boston, MA (Chair)

Irwin Brodsky, MD, Maine Medical Center Research Institute, Scarborough, ME

Lisa Ceglia, MD MS, Tufts Medical Center, Boston, MA

Chhavi Chadha, MD, HealthPartners Research Foundation, Minneapolis, MN

Ranee Chatterjee, MD MPH, Duke University Medical Center, Durham, NC

Bess Dawson-Hughes, MD, Jean Mayer USDA Human Nutrition Research Center on Aging at Tufts University, Boston, MA

Cyrus Desouza, MBBS, Omaha VA Medical Center, University of Nebraska Medical Center, Omaha, NE

Rowena Dolor, MD MHS, Duke University Medical Center, Durham, NC

John Foreyt, PhD, Baylor College of Medicine, Houston, TX

Adline Ghazi, MD, MedStar Good Samaritan Hospital, Baltimore, MD

Daniel S. Hsia, MD, Pennington Biomedical Research Center, Baton Rouge, LA

Karen C. Johnson, MD MPH, University of Tennessee Health Science Center, Memphis, TN

Sangeeta R. Kashyap, MD, Cleveland Clinic, Cleveland, OH

Sun Kim, MD, Stanford University Medical Center, Stanford, CA

Erin S. LeBlanc, MD MPH, Kaiser Permanente Center for Health Research NW, Portland, OR

Michael R. Lewis, MD MBA, Banner MD Anderson Cancer Center, Gilbert, AZ

Emilia Liao, MD, Northwell Health Lenox Hill Hospital, New York, NY

Saul Malozowski, MD PhD, National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda, MD

Lisa M. Neff, MD, Chicago, IL

Patrick O'Neil, PhD, Medical University of South Carolina, Charleston, SC

Jean Park, MD, MedStar Health Research Institute, Hyattsville, MD

Anne Peters, MD, Keck School of Medicine of the University of Southern California, Los Angeles, CA

Lawrence S. Phillips, MD, Atlanta VA Medical Center, Decatur, GA and Emory University School of Medicine, Atlanta, GA



Richard Pratley, MD, AdventHealth Translational Research Institute for Metabolism and Diabetes, Orlando, FL

Philip Raskin, MD, University of Texas Southwestern Medical Center, Dallas, TX

Neda Rasouli, MD, University of Colorado, School of Medicine and VA Eastern Colorado Health Care System, Aurora, CO

David Robbins, MD, University of Kansas Medical Center, Kansas City, KS

Clifford Rosen, MD, Maine Medical Center Research Institute, Scarborough, ME

Past Steering Committee members

Vanita R. Aroda, MD, Brigham and Women's Hospital, Boston, MA

Patricia Sheehan, RN MPH MS, Spaulding Rehabilitation Network, Boston, MA

Myrlene A. Staten, MD, National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda, MD

James H. Ware, PhD, Harvard T.H. Chan School of Public Health, Boston, MA (deceased)


Advisor

William C. Knowler, MD DrPH, National Institute of Diabetes and Digestive and Kidney Diseases, Phoenix, AZ




Vitamin **D** and
type **2** diabetes

Supplemental Figure S1: Consort diagram






Supplemental Figure S2: AUC_{glu} quintiles of screened population with mean fasting, 30-minute and 2-hour glucose levels




Quintiles were derived for the cross-sectional screening population (all races combined). Mean fasting, 30-minute and 2-hour levels in each quintile are shown in bold at the bottom of the graph. Total AUC_{glu} range (min-max) for Q1=9855 – 15480; Q2=15495 – 17205; Q3=17220 – 18795; Q4=18810 – 20820; Q5=20835 – 31245.

D_{2d} Vitamin **D** and type **2** diabetes



Original





Vitamin **D** and
type **2** diabetes

Supplemental Table S1. Baseline characteristics of screening cohort by race (cross-sectional analyses)

	N	Self-Identified Race			White vs Black p-value	White vs Asian p-value
		Black	White	Asian		
Characteristics		N=1016	N=2658	N=193	p-value	p-value
Age, years	3867	56.3 ± 9.7	61.2 ± 9.8	54.4 ± 11.1	<.0001	<.0001
Women, no. (%)	3867	504 (49.6)	1194 (44.9)	72 (37.3)	0.011	0.040
Body-mass index, kg/m ²	3866	32.5 ± 4.5	31.9 ± 4.4	28.1 ± 3.8	0.0005	<.0001
Body-mass index ≥30 kg/m ² , no. (%)	3867	703 (69.2)	1655 (62.3)	53 (27.5)	<.0001	<.0001
Hemoglobin, g/dL	3862	13.6 ± 1.3	14.4 ± 1.3	14.2 ± 1.2	<.0001	0.049
Presence of hemoglobin trait, no (%)						
C trait	3867	15 (1.5)	0 (0.0)	0 (0.0)	<.0001	NA
S trait	3867	58 (5.7)	1 (0.0)	0 (0.0)	<.0001	0.788
Serum 25(OH)D, ng/mL	3853	24.3 ± 10.7	29.8 ± 9.7	25.7 ± 9.6	<.0001	<.0001
Alcohol use, g/day, median (IQR)	3733	1.3 (0-6.5)	2.5 (0.2-13.1)	1.0 (0-4.3)	<.0001	<.0001
Smoking, no. (%)	3867				<.0001	0.088
Never/Former		905 (89.1)	2497 (93.9)	182 (94.3)		
Current		98 (9.6)	149 (5.6)	8 (4.1)		


**Vitamin D and
type 2 diabetes**

Unknown or Not reported		13 (1.3)	12 (0.5)	3 (1.6)		
Annual household income, no (%)	3867				<.0001	<.0001
Less than \$75,000		559 (55.0)	1184 (44.5)	45 (23.3)		
\$75,000 or more		270 (26.6)	1092 (41.1)	95 (49.2)		
Unknown or Not reported		187 (18.4)	382 (14.4)	53 (27.5)		
Education, no (%)	3866				<.0001	<.0001
No high school or post high school education		204 (20.1)	392 (14.7)	10 (5.2)		
Some post high school education		417 (41.0)	851 (32.0)	19 (9.9)		
Bachelor's degree		217 (21.4)	718 (27.0)	69 (35.9)		
Graduate or professional degree		175 (17.2)	679 (25.5)	90 (46.9)		
Unknown or Not reported		3 (0.3)	18 (0.7)	4 (2.1)		
Physical activity, total MET hour/week, median (IQR)	3767	63.0 (24.9-170.7)	56.8 (26.3-121.5)	43.2 (20.4-88.0)	0.0088	0.003
Dietary glycemic index, median (IQR)	3733	1024.5 (684.7-1539.6)	977.2 (715.9-1341.3)	981.8 (747.5-1392.4)	0.1442	0.494
Prediabetes/Diabetes status by test, no (%)						
A1c in prediabetes range	3859	903 (89.1)	2055 (77.5)	163 (84.9)	<.0001	0.016


**Vitamin D and
type 2 diabetes**

FPG in prediabetes range	3856	620 (61.1)	1994 (75.3)	142 (73.6)	<.0001	0.598
2hPG in prediabetes range	3849	319 (31.6)	935 (35.3)	73 (37.8)	0.0326	0.486
A1c in diabetes range	3859	17 (1.7)	30 (1.1)	4 (2.1)	0.1888	0.241
FPG in diabetes range	3856	23 (2.3)	136 (5.1)	2 (1.0)	0.0001	0.011
2hPG in diabetes range	3849	60 (5.9)	271 (10.2)	20 (10.4)	<.0001	0.957
No abnormal values	3851	259 (25.6)	355 (13.4)	27 (14.1)	<.0001	0.797



Vitamin **D** and
type **2** diabetes

Supplemental Table S2: Baseline characteristics of screening cohort by ethnicity (cross-sectional analyses)

	N	Ethnicity		Standardized mean difference (SMD)*
		Hispanic	Non-Hispanic	
		N=323	N=3544	
Characteristics				
Age, years	3867	54.3 ± 9.6	60.1 ± 10.1	<.0001
Women, no. (%)	3867	204 (63.2)	1566 (44.2)	<.0001
Body-mass index, kg/m ²	3866	32.4 ± 4.8	31.8 ± 4.5	0.0358
Body-mass index ≥30 kg/m ² , no. (%)	3867	212 (65.6)	2199 (62.0)	0.2029
Hemoglobin, g/dL	3862	13.9 ± 1.3	14.2 ± 1.3	<.0001
Presence of hemoglobin trait, no (%)				
C trait	3867	1 (0.3)	14 (0.4)	0.8131
S trait	3867	1 (0.3)	58 (1.6)	0.0625
Serum 25(OH)D, ng/mL	3853	26.4 ± 8.9	28.3 ± 10.4	0.0014
Alcohol use, g/day, median (IQR)	3733	0.8 (0-5.8)	2.2 (0.2-12.0)	<.0001
Smoking, no (%)	3867			0.6225
Never/Former		302 (93.5)	3282 (92.6)	
Current		20 (6.2)	235 (6.6)	
Unknown or Not reported		1 (0.3)	27 (0.8)	
Annual household income, no (%)	3867			<.0001
Less than \$75,000		210 (65.0)	1578 (44.5)	
\$75,000 or more		56 (17.3)	1401 (39.5)	
Unknown or Not reported		57 (17.6)	565 (15.9)	
Education, no (%)	3866			<.0001


**Vitamin D and
type 2 diabetes**

No high school or post high school education		170 (52.6)	436 (12.3)	
Some post high school education		76 (23.5)	1211 (34.2)	
Bachelor's degree		47 (14.6)	957 (27.0)	
Graduate or professional degree		28 (8.7)	916 (25.9)	
Unknown or Not reported		2 (0.6)	23 (0.6)	
Physical activity, total MET hour/week, median (IQR)	3767	79.0 (34.7-177.5)	56.0 (25.1-124.2)	<.0001
Dietary glycemic index, median (IQR)	3733	1065.6 (744.4-1612.3)	979.1 (708.1-1367.1)	0.0075
Glycemic categories, no. (%)	3851			0.7314
IGT + iA1c + IFG all in prediabetes range		63 (19.7)	788 (22.3)	
IGT + IFG in prediabetes range (A1c normal)		15 (4.7)	136 (3.9)	
IGT + iA1c in prediabetes range (FPG normal)		18 (5.6)	208 (5.9)	
IFG + iA1c in prediabetes range (2hPG normal)		97 (30.3)	1078 (30.5)	
Only one test in prediabetes range		28 (8.8)	318 (9.0)	
All 3 tests in normal range		63 (19.7)	578 (16.4)	
One test in diabetes range		36 (11.3)	425 (12.0)	



Supplemental Table S3: Baseline characteristics of enrolled cohort by race (longitudinal analyses)

	N	Race			p-value	
		White	Black	Asian	White vs Black	White vs Asian
		N=1613	N=616	N=130		
Characteristics						
Age, years	2359	61.7 ± 9.5	56.8 ± 9.5	55.2 ± 11.1	<.0001	<.0001
Women, no. (%)	2359	713 (44.2)	298 (48.4)	46 (35.4)	0.0767	0.0511
Body-mass index, kg/m ²	2359	32.1 ± 4.4	32.7 ± 4.5	28.2 ± 3.8	0.0045	<.0001
Body-mass index ≥30 kg/m ² , no. (%)	2359	1046 (64.8)	434 (70.5)	39 (30.0)	0.0122	<.0001
Hemoglobin, g/dL	2358	14.4 ± 1.2	13.7 ± 1.3	14.2 ± 1.3	<.0001	0.0729
Presence of hemoglobin trait, no (%)						
C trait	2359	0 (0.0)	10 (1.6)	0 (0.0)	<.0001	NA
S trait	2359	0 (0.0)	26 (4.2)	0 (0.0)	<.0001	NA
Serum 25(OH)D, ng/mL	2358	29.6 ± 9.5	24.2 ± 10.9	25.9 ± 10.2	<.0001	<.0001
Alcohol use, g/day, median (IQR)	2322	2.5 (0.2-13.1)	1.4 (0-6.7)	1 (0-5)	<.0001	0.0025
Smoking, no. (%)	2359				0.0007	0.0184
Never/Former		1517 (94.0)	552 (89.6)	122 (93.8)		
Current		89 (5.5)	56 (9.1)	5 (3.8)		
Unknown or Not reported		7 (0.4)	8 (1.3)	3 (2.3)		
Annual household income, no (%)	2359				<.0001	<.0001
Less than \$75,000		726 (45.0)	345 (56.0)	27 (20.8)		


**Vitamin D and
type 2 diabetes**

\$75,000 or more		662 (41.0)	164 (26.6)	61 (46.9)		
Unknown or Not reported		225 (13.9)	107 (17.4)	42 (32.3)		
Education, no (%)	2359				<.0001	<.0001
No high school or post high school education		230 (14.3)	137 (22.2)	7 (5.4)		
Some post high school education		514 (31.9)	248 (40.3)	13 (10.0)		
Bachelor's degree		456 (28.3)	126 (20.5)	48 (36.9)		
Graduate or professional degree		404 (25.0)	104 (16.9)	60 (46.2)		
Unknown or Not reported		9 (0.6)	1 (0.2)	2 (1.5)		
Physical activity, total MET hour/week, median (IQR)	2318	54.6 (25.8-119.5)	63 (26.6-169.5)	39.9 (18.2-75.6)	0.0094	0.0018
Dietary glycemic index, median (IQR)	2322	977 (714.5-1331)	1035.3 (691.2-1573)	949.2 (762.7-1312.4)	0.0531	0.8491
Glycemic categories, no. (%)	2359				<.0001	0.3037
IGT + iA1c + IFG all in prediabetes range		602 (37.3)	185 (30.0)	46 (35.4)		
IGT + IFG in prediabetes range (A1c normal)		126 (7.8)	17 (2.8)	6 (4.6)		
IGT + iA1c in prediabetes range (FPG normal)		119 (7.4)	91 (14.8)	14 (10.8)		
IFG + iA1c in prediabetes range (2hPG normal)		766 (47.5)	323 (52.4)	64 (49.2)		



Vitamin **D** and
type **2** diabetes

Supplemental Table S4: Baseline characteristics of enrolled cohort by ethnicity

	N	Ethnicity		Hispanic vs non-Hispanic p-value
		Hispanic	non-Hispanic	
		N=190	N=2169	
Characteristics				
Age, years	2359	54.6 ± 9.8	60.6 ± 9.8	<.0001
Women, no. (%)	2359	118 (62.1)	939 (43.3)	<.0001
Body-mass index, kg/m ²	2359	32.5 ± 4.4	32.0 ± 4.5	0.1909
Body-mass index ≥30 kg/m ² , no. (%)	2359	127 (66.8)	1392 (64.2)	0.462
Hemoglobin, g/dL	2358	13.9 ± 1.3	14.2 ± 1.3	0.0008
Presence of hemoglobin trait, no (%)				
C trait	2359	1 (0.5)	9 (0.4)	0.8207
S trait	2359	0 (0.0)	26 (1.2)	0.1291
Serum 25(OH)D, ng/mL	2358	25.9 ± 8.1	28.2 ± 10.4	0.0033
Alcohol use, g/day, median (IQR)	2322	0.8 (0-6.3)	2.2 (0.2-12.1)	<.0001
Smoking, no. (%)	2359			0.4228
Never/Former		179 (94.2)	2012 (92.8)	
Current		11 (5.8)	139 (6.4)	
Unknown or Not reported		0 (0.0)	18 (0.8)	
Annual household income, no (%)	2359			<.0001
Less than \$75,000		123 (64.7)	975 (45.0)	
\$75,000 or more		35 (18.4)	852 (39.3)	


**Vitamin D and
type 2 diabetes**

Unknown or Not reported		32 (16.8)	342 (15.8)	
Education, no (%)	2359			<.0001
No high school or post high school education		95 (50.0)	279 (12.9)	
Some post high school education		47 (24.7)	728 (33.6)	
Bachelor's degree		32 (16.8)	598 (27.6)	
Graduate or professional degree		15 (7.9)	553 (25.5)	
Unknown or Not reported		1 (0.5)	11 (0.5)	
Physical activity, total MET hour/week, median (IQR)	2318	67 (32.6-182.6)	54.1 (25-121.8)	0.0021
Dietary glycemic index, median (IQR)	2322	1053.5 (711.6-1471.6)	981 (711.3-1377.2)	0.3462
Glycemic categories, no (%)	2359			0.6749
IGT + iA1c + IFG ⁴ all in prediabetes range		61 (32.1)	772 (35.6)	
IGT + IFG in prediabetes range (A1c normal)		15 (7.9)	134 (6.2)	
IGT + iA1c in prediabetes range (FPG normal)		18 (9.5)	206 (9.5)	
IFG + iA1c in prediabetes range (2hPG normal)		96 (50.5)	1057 (48.7)	



Vitamin **D** and
type **2** diabetes

Supplemental Table S5: A1c level by AUC-glucose quintiles in screened population by ethnicity

AUC _{glu} concentration quintiles	Hispanic		Non-Hispanic	
	Observed A1c	Adjusted A1c [†]	Observed A1c	Adjusted A1c [†]
Mean(range)				
Low-1	5.79 (5.73, 5.85)	5.81 (5.74, 5.88)	5.81 (5.79, 5.83)	5.82 (5.80, 5.84)
2	5.86 (5.79, 5.92)	5.86 (5.79, 5.93)	5.84 (5.82, 5.86)	5.84 (5.82, 5.86)
3	5.89 (5.83, 5.95)	5.89 (5.83, 5.95)	5.86 (5.84, 5.88)	5.86 (5.84, 5.88)
4	5.83 (5.77, 5.89) [‡]	5.84 (5.77, 5.90)	5.90 (5.88, 5.92)	5.89 (5.87, 5.91)
High-5	6.05 (5.98, 6.12) [‡]	6.05 (5.98, 6.12) [‡]	5.97 (5.95, 5.99)	5.96 (5.95, 5.98)

*Quintiles were derived for the screening population (all races combined)

[†]Estimated linear regression for adjusted A1c = (race) + AUC quintile groups + (race)*AUC quintile groups + adjusted variables (age, gender, BMI, age*gender*BMI, alcohol use, education, income, presence of hemoglobinopathy, smoking status, physical activity, dietary glycemic index, and 25[OH]D)

[‡]p<0.05 for Nonhispanic vs Hispanic participants



Supplemental Table S6: Change in observed A1c over time by baseline AUC-glucose quintile and race in enrolled cohort (longitudinal)

Race	Baseline AUC _{glu} quintile	Observed A1c (Mean, SD)	Change in A1c over time					Comparison of change in A1c in Black vs White participants				
			Baseline	Year1	Year2	Year3	Year4	Percent difference (95% CI)	P value	Percent difference (95% CI)	Unadjusted p-value	Adjusted p-value [†]
White	1-Low	N 257	237	207	107	34			ref	ref	ref	
			5.86 (0.16)	5.82 (0.24)	5.88 (0.24)	5.88 (0.22)	5.86 (0.33)	0.03 (-0.37, 0.43)				0.884
	2	N 312	284	243	132	48						
			5.89 (0.18)	5.9 (0.23)	5.94 (0.26)	5.93 (0.25)	5.96 (0.24)	0.85 (0.48, 1.21)				<.001
	3	N 314	286	238	129	54						
			5.87 (0.21)	5.86 (0.24)	5.92 (0.25)	5.92 (0.25)	5.96 (0.27)	0.69 (0.33, 1.06)				<.001
Black	4	N 344	311	222	114	39						
			5.88 (0.22)	5.9 (0.29)	5.9 (0.27)	5.93 (0.31)	5.93 (0.28)	0.85 (0.49, 1.21)	<.001			
	5-High	N 372	317	219	109	37						
			5.91 (0.23)	5.92 (0.27)	5.97 (0.32)	5.91 (0.28)	5.93 (0.27)	1.19 (0.83, 1.55)	<.001			
	1-Low	N 195	168	139	90	41						
			5.96 (0.18)	5.94 (0.21)	5.96 (0.24)	5.99 (0.27)	5.92 (0.32)	0.76 (0.29, 1.23)	0.002	0.73 (0.11, 1.35)	0.022	0.057
Black	2	N 133	113	90	58	24						
			6.04 (0.19)	6.04 (0.24)	6.08 (0.23)	6.11 (0.25)	6.08 (0.19)	1.81 (1.23, 2.40)	<.001	0.97 (0.27, 1.66)	0.006	0.027
	3	N 122	103	80	45	22						
			5.98 (0.19)	5.99 (0.21)	6.04 (0.23)	6.05 (0.21)	6.15 (0.19)	1.26 (0.65, 1.88)	<.001	0.57 (-0.15, 1.28)	0.121	0.243
	4	N 93	81	61	42	12						
			6.02 (0.23)	5.98 (0.3)	6 (0.26)	6.04 (0.27)	5.96 (0.38)	0.88 (0.18, 1.57)	0.013	0.03 (-0.76, 0.81)	0.945	0.737
Black	5-High	N 66	54	37	20	8						
			6.02 (0.25)	5.93 (0.3)	6.11 (0.4)	6.04 (0.21)	6.11 (0.32)	1.19 (0.33, 2.05)	0.007	0.00 (-0.93, 0.94)	0.998	0.947

* Percent difference in A1c refers to least square mean difference in the longitudinal values over time using linear mixed model.

[†]Adjusted for baseline variables: race, AUC quintile groups, interaction term with race and AUC quintile groups, baseline A1c, age, gender, BMI, interaction terms with age, gender, and BMI, FFQ alcohol, education, income, presence of hemoglobinopathy, smoking status, physical activity, dietary glycemic index, 25(OH)D, and randomization assignment.